



Algebra 2 3.5

Analyze data in matrices

Perform algebraic operations on matrices

distributive property  $3 \times 5$   
matrix (matrices pl.)  $r \times c$   
row - - .  
column .  
element \*  
corresponding elements  
dimension .  
scale factor .  
scalar .

$$\begin{bmatrix} 5 & 0 & 6 & 7 & -5 \\ 1 & -3 & 2 & 1 & 5 \\ 9 & 0 & 2 & 1 & 5 \end{bmatrix}$$

$$\begin{bmatrix} x & a \\ n & s \end{bmatrix}$$

$$2 \times 2$$

$$\begin{bmatrix} 1 & 3 \\ 6 & 4 \end{bmatrix}$$

$$2 \times 2$$

whiteboards


Sample matrix

$$\begin{bmatrix} 1 & 5 & 7 \\ 3 & 8 & -6 \end{bmatrix} - \begin{bmatrix} 0 & -3 & 5 \\ 8 & 1 & 7 \end{bmatrix} = \begin{bmatrix} 1 & 8 & 2 \\ -5 & 7 & -13 \end{bmatrix}$$

$2 \times 3$                        $2 \times 3$

$$\begin{bmatrix} 1 & 5 & 7 \\ 3 & 8 & -6 \end{bmatrix} + \begin{bmatrix} 1 & 6 \\ 2 & 5 \end{bmatrix} = NP$$

**2 Algebraic Operations** Several algebraic operations can be performed on data that are organized in matrices. Matrices can be added or subtracted if and only if they have the same dimensions.

 **KeyConcept** Adding and Subtracting Matrices

**Words** To add or subtract two matrices with the same dimensions, add or subtract their corresponding elements.

**Symbols**

$$\mathbf{A} + \mathbf{B} = \mathbf{A} + \mathbf{B}$$
$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} + \begin{bmatrix} e & f \\ g & h \end{bmatrix} = \begin{bmatrix} \underline{a+e} & \underline{b+f} \\ c+g & d+h \end{bmatrix}$$
$$\mathbf{A} - \mathbf{B} = \mathbf{A} - \mathbf{B}$$
$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} - \begin{bmatrix} e & f \\ g & h \end{bmatrix} = \begin{bmatrix} a-e & b-f \\ c-g & d-h \end{bmatrix}$$

**Example**

$$\begin{bmatrix} 3 & -5 \\ 1 & 7 \end{bmatrix} + \begin{bmatrix} 2 & 0 \\ -9 & 10 \end{bmatrix} = \begin{bmatrix} 3+2 & -5+0 \\ 1+(-9) & 7+10 \end{bmatrix}$$

(if possible)



**Example 2** Add and Subtract Matrices

Find each of the following for  $A = \begin{bmatrix} 16 & 2 \\ -9 & 8 \end{bmatrix}$ ,  $B = \begin{bmatrix} -4 & -1 \\ -3 & -7 \end{bmatrix}$ , and  $C = \begin{bmatrix} 8 \\ 6 \end{bmatrix}$ .

a.  $A + B = \begin{bmatrix} 12 & 1 \\ -12 & 1 \end{bmatrix}$        $2 \times 2$        $2 \times 2$        $2 \times 1$

b.  $B - C = \begin{bmatrix} \phantom{0} & \phantom{0} \\ \phantom{0} & \phantom{0} \end{bmatrix}$  NP

c.  $B - A = \begin{bmatrix} -20 & -3 \\ 6 & -15 \end{bmatrix}$

SAME dimensions!

whiteboards

### Guided Practice

$$2A. \begin{bmatrix} -3 & 4 \\ -9 & -5 \end{bmatrix} - \begin{bmatrix} -4 & 12 \\ 8 & -7 \end{bmatrix}$$

$$2C. \begin{bmatrix} 8 & -3 \\ -2 & 0 \\ 1 & 7 \end{bmatrix} - \begin{bmatrix} 5 & 1 & -4 & 2 \\ 10 & -6 & 9 & 0 \end{bmatrix}$$

$$3+3+3+3+3 = 3 \cdot 5 =$$

Scalar  
mult.

$$A + A + A + A + A = 5A$$

$$\begin{bmatrix} 2 & 1 \\ 3 & 0 \end{bmatrix}$$

$$\begin{bmatrix} 10 & 5 \\ 15 & 0 \end{bmatrix}$$

**Example 3** Multiply a Matrix by a Scalar

If  $R = \begin{bmatrix} -12 & 8 & 6 \\ -16 & 4 & 19 \end{bmatrix}$ , find  $5R$ .

$$R + R + R + R + R$$

$$\begin{bmatrix} -60 & 40 & 30 \\ -80 & 20 & 95 \end{bmatrix}$$

Scalar...scale factor



### Guided Practice

3. If  $T = \begin{bmatrix} 8 & 0 & 3 & -2 \\ -1 & -4 & -2 & 9 \end{bmatrix}$ , find  $-4T$ .

### KeyConcept Properties of Matrix Operations

For any matrices  $A$ ,  $B$ , and  $C$  for which the matrix sum and product are defined and any scalar  $k$ , the following properties are true.

Commutative Property of Addition  $A + B = B + A$

Associative Property of Addition  $(A + B) + C = A + (B + C)$

Left Scalar Distributive Property  $k(A + B) = kA + kB$

Right Scalar Distributive Property  $(A + B)k = kA + kB$

scalar multiplication is not the same as matrix multiplication...

Order of operations.... GEMA

### Example 4 Multi-Step Operations

If  $A = \begin{bmatrix} -9 & 12 \\ 2 & -6 \end{bmatrix}$  and  $B = \begin{bmatrix} -4 & -8 \\ 2 & -3 \end{bmatrix}$ , find  $-4B + 3A$ .

$$\begin{array}{c} -4B \\ \left[ \begin{array}{cc} 16 & 32 \\ -8 & +12 \end{array} \right] \end{array} + \begin{array}{c} -3A \\ \left[ \begin{array}{cc} 27 & -36 \\ -6 & 18 \end{array} \right] \end{array}$$

### Guided Practice

4. If  $A = \begin{bmatrix} -5 & 3 \\ 6 & -8 \\ 2 & 9 \end{bmatrix}$  and  $B = \begin{bmatrix} 12 & 5 \\ 5 & -4 \\ 4 & -7 \end{bmatrix}$ , find  $-6B + 7A$ .

$$\begin{matrix} -6B \\ \left[ \begin{array}{cc} & \end{array} \right] \end{matrix} + \begin{matrix} 7A \\ \left[ \begin{array}{cc} & \end{array} \right] \end{matrix} = \begin{bmatrix} -107 & -9 \\ 12 & -32 \\ -10 & 105 \end{bmatrix}$$



p. 174 wages Sales 5-day

	wages	Sales
Entry	900	145,000
Assist	2400	225,000
Assoc	2700	290,000

$$5 \begin{bmatrix} 900 & 145,000 \\ 2400 & 225,000 \\ 2700 & 290,000 \end{bmatrix} =$$

3.5 13-270 37-510

P175

$$700 \times .06 = 42$$

$$700 + 42 =$$

$$700 \times 1.06$$