

Trig 8.2

$\langle 3, 5 \rangle$

Find ordered pairs that represent vectors
Add, subtract, multiply and divide vectors

vector

resultant

vertical

horizontal


component

unit vector

cosine (x adjacent)

sine (y opposite)


$\tan^{-1} () \rightarrow \text{angles}$




Is a geometric method adequate to combine vectors?

Is it precise enough?

Is it user friendly and convenient?



Standard position
magnitude...direction... (same?)



Ordered pairs or vectors?

Some books use $\langle \quad \rangle$ to represent vectors
<component form> $\langle x, y \rangle$


"brackets"

- 1 Write the ordered pair that represents the vector from $X(-3, 5)$ to $Y(4, -2)$.
Then find the magnitude of \overline{XY} .

$$\langle 7, -7 \rangle$$

$$4 - (-3)$$
$$-2 - 5$$

$$7^2 + 7^2 =$$
$$49 + 49 = h^2$$
$$\sqrt{98} = h$$
$$\ominus$$
$$7\sqrt{2} = h$$
$$\approx 9.9$$



(When it's in component form...)


$\langle x, y \rangle$

Write the ordered pair that represents \overline{MP} . Then find the magnitude of \overline{MP} .

4. $M(2, -1), P(-3, 4)$

5. $M(5, 6), P(0, 5)$

6. $M(-19, 4), P(4, 0)$



Much easier to add & subtract when in **component** form!
 (How can you tell?)

Vector Operations

The following operations are defined for $\vec{a} \langle a_1, a_2 \rangle$, $\vec{b} \langle b_1, b_2 \rangle$, and any real number k .


Addition: $\vec{a} + \vec{b} = \langle a_1, a_2 \rangle + \langle b_1, b_2 \rangle = \langle a_1 + b_1, a_2 + b_2 \rangle$

Subtraction: $\vec{a} - \vec{b} = \langle a_1, a_2 \rangle - \langle b_1, b_2 \rangle = \langle a_1 - b_1, a_2 - b_2 \rangle$

Scalar multiplication: $k\vec{a} = k\langle a_1, a_2 \rangle = \langle ka_1, ka_2 \rangle$

Example 2 Let $\vec{m} = \langle 5, -7 \rangle$, $\vec{n} = \langle 0, 4 \rangle$, and $\vec{p} = \langle -1, 3 \rangle$. Find each of the following.

b. $\vec{m} + \vec{n}$ SMATO



$$\begin{array}{r} \langle 5, -7 \rangle \\ + \langle 0, 4 \rangle \\ \hline \langle 5, -3 \rangle \end{array}$$


d. $2\vec{m} + 3\vec{n} + \vec{p}$

$$\begin{array}{r} \langle 10, -14 \rangle \\ + \langle 0, 12 \rangle \\ + \langle -1, -3 \rangle \\ \hline \langle 9, -5 \rangle \end{array}$$

24° S of E

$$\langle 11, -5 \rangle$$

336° from horz



$\tan^{-1} \frac{1}{5}$
 $\theta = 24^\circ$

66° E of S

(Is it already in component form?)

Find an ordered pair to represent \vec{t} in each equation if $\vec{u} = \langle -1, 4 \rangle$ and $\vec{v} = \langle 3, -2 \rangle$.

7. $\vec{t} = \vec{u} + \vec{v}$

8. $\vec{t} = \frac{1}{2}\vec{u} - \vec{v}$


$$\vec{t} = \langle -1, 4 \rangle + \langle 3, -2 \rangle$$

$$\vec{t} = \langle 2, 2 \rangle$$

$$\langle 2, 3 \rangle$$

A vector that has a magnitude of one unit is called a **unit vector**. A unit vector in the direction of the positive x -axis is represented by \vec{i} , and a unit vector in the direction of the positive y -axis is represented by \vec{j} . So, $\vec{i} = \langle 1, 0 \rangle$ and $\vec{j} = \langle 0, 1 \rangle$.

$$2\vec{i} + 3\vec{j}$$




So how do we know whether i represents a unit vector or $\sqrt{-1}$?

Components $\langle x, y \rangle$ unit vectors i & j

4 Write \vec{AB} as the sum of unit vectors for $A(4, -1)$ and $B(6, 2)$.

$\vec{a} + \vec{b}$
 $\star 2\vec{i} + 3\vec{j}$

x, y
 h, v
 \vec{i}
 \vec{j}




i & j

Find the magnitude of each vector. Then write each vector as the sum of unit vectors.

11. $\langle 8, -6 \rangle$


$$8\vec{i} - 6\vec{j}$$

$$10$$



12. $\langle -7, -5 \rangle$

$$-7\vec{i} - 5\vec{j}$$




Find the magnitude of each vector. Then write each vector as the sum of unit vectors.

36. $\langle 3, 4 \rangle$

37. $\langle 2, -3 \rangle$

38. $\langle -6, -11 \rangle$



Algebraic Vectors




EMERGENCY MEDICINE

Paramedics Paquita Gonzalez and Trevor


Howard are moving a person on a stretcher. Ms. Gonzalez is pushing the stretcher with a force of 135 newtons at 58° with the horizontal, while Mr. Howard is pulling the stretcher with a force of 214 newtons at 43° with the horizontal. What is the magnitude of the force exerted on the stretcher? *This problem will be solved in Example 3.*



$$\begin{array}{r} \langle -71.5, 114.5 \rangle \\ \langle 156.5, 145.9 \rangle \\ \hline \langle 85, 260.4 \rangle \end{array}$$




$$\begin{array}{l} 135 \\ \swarrow 58^\circ \\ x = -71.5 \\ y = 114.5 \end{array}$$



- components
- combine
- magnitude

magnitude & direction

13. **Construction** The Walker family is building a cabin for vacationing. Mr. Walker and his son Terrell have erected a scaffold to stand on while they build the walls of the cabin. As they stand on the scaffold Terrell pulls on a rope attached to a support beam with a force of 400 newtons (N) at an angle of 65° with the horizontal. Mr. Walker pulls with a force of 600 newtons at an angle of 110° with the horizontal. What is the magnitude of the combined force they exert on the log?



and direction

